

Number concentration and modal Structure of indoor/outdoor fine particles in four European Cities

M. Lazaridis¹, K. Eleftheriadis², V. Ždímal³, J. Schwarz³, Z. Wagner³, J. Ondracek³, Y. Drossinos⁴, T. Glytsos¹, S. Vratolis², K. Torseth⁴ and J. Smolik³

¹Department of Environmental Engineering, Technical University of Crete, Chania 73100, Greece,

²Institute of Nuclear Technology & Radiation Protection, N.C.S.R. “Demokritos”, 15310 Ag. Paraskevi, Attiki, Greece,

³Institute of Chemical Process Fundamentals, Laboratory of Aerosol Chemistry and Physics, Rozvojová 135, CZ-16502 Praha 6, Prague, Czech Republic,

⁴European Commission, Joint Research Centre, I-21020 Ispra (Va), Italy.

⁵Norwegian Institute for Air Research (NILU), Instituttveien 18, N-2020 Kjeller, Norway

Keywords: Indoor number concentration, indoor sources, modal structure.

Presenting author email: thodoros.glytsos@enveng.tuc.gr

Indoor/outdoor measurements campaigns have been performed in naturally ventilated houses and offices in four European cities in order to investigate the relations between indoor and outdoor particle number concentrations and aerosol modal structures. The measurements sites were located in Oslo (Norway), Prague (Czech Republic), Milan (Italy) and Athens (Greece). A number of ordinary indoor activities (cooking, smoking, candle burning, etc.) were performed during the measurement's periods with the aim to quantify the contribution of the emitted particles from indoor sources and also the contribution of outdoor particles to the change to the indoor aerosol size spectra.

The size and time resolved particle concentrations in the size range 10 nm – 600 nm were measured by scanning mobility particle sizers (SMPS). The results were analysed, and the modal structure of the measured particle number size distribution data along with the number concentration in each of the modes comprising the aerosol size distribution were computed.

The analysis of the measurements showed that in absence of indoor sources, indoor number concentration followed the variations of the outdoor concentration. Figure 1 presents the correlation between diurnal median values of indoor and outdoor number concentrations for the measurements conducted in Prague. Indoor concentration was highly correlated to the outdoor concentration when the apartment was unoccupied, but poor correlation was observed during periods with indoor activities.




Figure 1. Correlation of indoor and outdoor diurnal median number concentrations for the measurements conducted in Prague (Czech Republic).

Emissions from indoor sources resulted in elevated indoor number concentrations and I/O ratios. Furthermore, the aerosol spectra presented different characteristics during periods with indoor activities as it is depicted in Table 1. Outdoor size distribution was similar to the indoor size distribution, which suggests that in naturally ventilated buildings, the indoor size spectra is formed according to the penetration efficiency of the outdoor particles to the indoor environment and to the emission characteristics of indoor sources.

Table 1a,b: Median values of GMDs, sorted according to time period without indoor activities(a) and with indoor activities (b) in the measurements sites of Oslo, Prague and Milan. The numbers 1, 2 refer to the corresponding mode (1 for 1st mode, etc.)

Median values for time periods without indoor activities			
	GMD [nm]		
	1	2	
Unimodal	Prague	120	
	Oslo	57	
	Milan	74	
	Athens		
Bimodal	Prague	34	132
	Oslo	23	91
	Milan	41	128

(a)

Median values for periods with indoor activities			
	GMD [nm]		
	1	2	
Unimodal	Prague	85	
	Oslo	85	
	Milan	129	
	Athens		
Bimodal	Prague	45	182
	Oslo	20	101
	Milan	48	150

(b)

The research leading to these results has received funding from the European Union Seventh Framework Programme HEXACOMM FP7/2007-2013 under grant agreement n° 315760.